



WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

PCT

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 16/00, 16/46, 19/00, A61K 48/00, 51/00, 121/00		A1	(43) International Publication Date: 10 August 1995 (10.08.95)
(11) International Publication Number: WO 95/21195			
(21) International Application Number: PCT/US95/01161		(81) Designated States: AU, CA, CN, FI, JP, KR, NO, NZ, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 6 February 1995 (06.02.95)		(30) Priority Data: 08/192,655 US	
(60) Parent Application or Grant (63) Related by Continuation US Filed on 08/192,655 (CIP) 7 February 1994 (07.02.94)		(71) Applicant (for all designated States except US): RESEARCH DEVELOPMENT FOUNDATION [US/US]; 402 North Di- vision Street, Carson City, NV 89703 (US).	
(72) Inventors; and (75) Inventors/Applicants (for US only): ROSENBLUM, Michael, G. [US/US]; 3127 Stonewall Drive, Sugar Land, TX 77479 (US), DONATO, Nicholas, J. [US/US]; 6206 Bank- side, Houston, TX 77096 (US).		(74) Agent: WEILLER, James, F.; Suite 1560, 1 Riverway, Houston, TX 77056 (US).	
(54) Title: NON-VIRAL VECTOR			
(57) Abstract <p>The present invention provides a non-viral vector, comprising a cell binding component having a biotin-binding element conjugated to a biotinylated moiety. Also, provided is a method of introducing genetic material inside a specific cell comprising the administration of the non-viral vector to a human, wherein said non-viral vector comprises a cell binding component having a biotin-binding element conjugated to a biotinylated moiety. In addition, there is provided a method of delivering a cytotoxic moiety to a cell comprising the administration of the non-viral vector to a human.</p>			

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

FOR THE PURPOSES OF INFORMATION ONLY

AT	Austria	GB	United Kingdom	MIR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland			SI	Slovenia
CI	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	LI	Liechtenstein	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali	UZ	Uzbekistan
FR	France	MN	Mongolia	VN	Viet Nam
GA	Gabon				

30 The present invention relates generally to the field of molecular biology and therapeutics. More specifically, the present invention relates to a novel non-viral vector for the delivery of genetic material capable of affecting molecular properties of mammalian cells utilizing viral, primarily retroviral, vectors. However, this mode of genetic therapy or delivery is influenced by several deficiencies and potentially hazardous conditions. Although retroviral vectors have been more common, adenoviral vectors are now being studied and both have some potential. Unfortunately, a viral vector has many disadvantages. First, the target cells, e.g., human cells, must be capable of interacting with viruses through expression of a specific cell surface element which may not be expressed on the cells or tissue of interest for delivery of genetic information. Second, the genetic material must be integrated and expressed in the target, e.g., human cell, which requires that target cells be actively dividing, a condition hindering the efficiency and homogeneity of this delivery system. Even if successfully integrated, the gene may be transcriptionally silent by host cell mechanisms. Third, the size of the genetic information allowable in this system is limited and must be engineered with great precision to ensure biologic activity. Fourth, the replication defective viruses must be utilized for gene therapy applications to reduce the risk of recombination with endogenous viruses which may form new infectious agents. Replication defective viruses may reduce

Field of the Invention

The present invention relates generally to the field of molecular biology and therapeutics. More specifically, the present invention relates to a novel non-viral vector for the delivery of genetic information to cells.

Description of the Related Art

BACKGROUND OF THE INVENTION

NON-VIRAL VECTOR

5 this hazard but do not eliminate it. Fifth, the replication defective viruses are by design not self-removing, requiring repetitive infection to achieve successful delivery of gene sequences to all cells.

The prior art is deficient in the absence of non-viral vectors. The present invention fulfills this longstanding need and desire in the art.

SUMMARY OF THE INVENTION

10 The present invention provides, *inter alia*, a novel non-viral system for delivery of genetic materials capable of modification of deleterious or undesirable phenotypic characteristics.

Thus, in one embodiment of the present invention there is provided a non-viral vector, comprising a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

20 In another embodiment of the present invention, there is provided a method of introducing genetic material inside a specific cell comprising the administration of the non-viral vector to a human, wherein said non-viral vector comprises a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

25 In yet another embodiment of the present invention, there is provided a method of delivering a cytotoxic moiety to a cell comprising the administration of the non-viral vector to a human.

30 Other and further aspects, features, and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention given for the purpose of disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

The following figures are provided to illustrate various aspects of the present invention. To that end, some of the figures are presented in schematic form and are not necessarily drawn to scale.

Figure 1 shows the separation of free SPDP from A108/SPDP on G-25 column.

Figure 2 shows the separation of free 2-IT from avidin/2-IT on G-25 column.

Figure 3 shows the separation of free avidin from A108-avidin conjugate on S-200 column (FPLC).

Figure 4 shows the separation of free antibody from A108-avidin conjugate on Con-A column.

Figure 5 shows a 7.5% SDS-PAGE mini-gel showing purification steps of A108-avidin conjugate.

Figure 6 shows the binding activity for assay of biotinylated gelonin.

Figure 7 shows the profile of G-75 (FPLC) for A108-avidin gelonin/biotin conjugate and the separation of free avidin gelonin and some avidin sub-units.

Figure 8 shows the Elisa binding activity assay for biotinylated gelonin and for A108-avidin biotinylated gelonin conjugate.

Figure 9 shows the cytotoxicity of A108-gelonin conjugate compared with the A108-avidin biotinylated gelonin conjugate on A431 cells.

Figure 10 shows the effect of incubation of cells with nucleic acid sequences directed against the EGF receptor gene promoter sequence (labeled anti-sense EGF)

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a non-viral vector, comprising a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

Generally, the biotin-binding element of the present invention is any that chemical that binds biotin and would be easily recognizable by a person having ordinary skill in this art. Preferably, the biotin-binding element is selected from the group consisting of avidin, streptavidin or analogues of avidin or streptavidin.

The cell binding element of the present invention may be one of several different embodiments. For example, the

cell binding element may be a monoclonal antibody. Monoclonal antibody useful in the compositions and methods of the present invention are those that specifically bind an antigen. Representative examples of antigens to which such antibodies would bind are epidermal growth factor receptor, c-erbB2 antigen, Lewis Y antigen, transferrin receptor, MDR1, MDR3, insulin receptor, CD45, CD33, GP240, GD2, GD3, fibroblast growth factor receptor, platelet derived growth factor receptor. Alternatively, the cell binding element is a ligand which specifically binds a cell surface receptor. Representative examples of ligands binding cell surface receptors include transforming growth factor-alpha, heregulin, fibroblast growth factor, platelet-derived growth factor receptor. Generally, the biotinylated moiety may be any compound which can be appropriately biotinylated and which is a chemical which one desires to specifically introduce inside a cell to exert a particular biological or pharmacological effect. Thus, the biotinylated moiety may be a protein or a nucleic acid.

Representative examples of proteins useful in the compositions and methods of the present invention are gelonin, ricin, saporin, abrin, diphtheria toxin, pseudomonas exotoxin, rayalase, superoxide dismutase, protein tyrosine phosphatase, protein phosphatase (PP-1 or PP-2), protein kinase A and protein kinase C.

Representative examples of nucleic acids are triple helix oligonucleotides, e.g., triplex EGF receptor oligonucleotides, anti-sense oligonucleotides, e.g., for EGF or myc, partial gene sequences, e.g., sequences encoding a single domain of a protein with several domains such as c-src or the EGF receptor and entire genes, i.e., taken from an integrative unit of the retroviral genome.

The present invention also provides a method of introducing genetic material inside a specific cell comprising

the administration of the non-viral vector to a human, wherein said non-viral vector comprises a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

5 The present invention also provides a method of delivering a cytotoxic moiety to a cell comprising the administration of the non-viral vector to a human.

The present invention involves compositions and methods permitting the introduction of nucleic acids into a specific subset of cells without using a viral infection or transfection component. Monoclonal antibodies directed against a cell-surface component are modified and utilized to carry, to the intracellular compartment, nucleic acids capable of modifying gene expression, specifically increasing or decreasing the level of protein expressed within target cells.

15 The present invention is applicable for anti-sense nucleic acid technology in humans.

Monoclonal antibodies directed against a cell surface component are modified with a biotin-binding moiety and nucleic acid sequences that bind genomic DNA or mRNA sequences such that gene expression is altered and are linked through avidin:biotin interaction using biotinylated derivatives of nucleic acid sequences. Through internalization of antibody:antigen complexes, active nucleic acid sequences are colocalized, increasing the intracellular content of these sequences. Use of the present invention: (1) increases intracellular content of gene expression modulating nucleic acid sequences by mechanisms other than those related to viral vectors or through simple or facilitated diffusion through the plasma membrane; (2) allows one to bypass the restrictions of small molecular size, i.e., of active nucleic acid sequences (-20-mers) by eliminating the need of simple diffusion through the plasma membrane as the mechanism of entry of these sequences into the cell; and (3) delivers sequences of interest to cells specifically or selectively through an antibody:antigen interaction rather

-6-

than through global delivery or viral infection to increase cellular content of active nucleic acid sequences.

In one embodiment of the present invention, one synthesizes anti-sense DNA against an oncogenic protein which spans nucleotides (10) upstream and downstream of the mRNA translation start codon. Then, one synthesizes DNA complementary to the first 5 nucleotides of the anti-sense DNA synthesized above. Then, one incorporates into that sequence a biotin-nucleotide moiety. The two strands are hybridized. Then, deliver via tumor targeting the MAb Avidin/Streptavidin to tumor.

The following examples are provided for the sole purpose of illustrating various embodiments of the present invention and are not meant to limit the present invention in any fashion.

EXAMPLE 1

Modification of Antibody A108

A108 recognizes the human receptor for epidermal growth factor. 10 mg of A108 in 2.2 ml of phosphate-buffered saline was added to a 12 x 75 mm glass tube. An aliquot of 9.35 μ l of antibody and a 2.5 fold molar excess of SPDP (N-succinimidyl 3-(2-pyridylidithio)propionate) from a stock of 3 mg/ml in dimethyl formamide was added slowly to the tube while vortexing. The mixture was vortexed every five minutes during a 30 minute incubation at room temperature.

Excess unreacted SPDP was removed from the sample by gel filtration chromatography on a column (1.5 x 37 cm) of Sephadex G-25 pre-equilibrated in 100 mM sodium phosphate buffer (pH 7.0) containing 0.5 mM EDTA. One ml fractions were collected on a Gilson fraction collector during buffer elution. Fractions were analyzed for protein content in a 96-well microtiter plate (Falcon) using the Bradford dye binding assay. Each well contained 120 μ l of PBS, 40 μ l of dye concentrate and 40 μ l of sample. Absorbance was read on a Biotek Microplate Autoreader at 540 nm. Fractions 30-38 were pooled and kept at 4°C.

35 groups, and incubation continued for 1 hour at room temperature. SPDP-modified A108 and 2-iminothiolan-modified
 30 The modified antibody and modified avidin fractions
 conjugation of antibody A108 and avidin

EXAMPLE 3

Figure 2.
 25 iminothiolane was recovered by gel filtration as shown in
 (1 ml each) 27-38 were pooled. Avidin modified with 2-
 read on a Biotek Microplate Autoreader at 540 nm. Fractions
 determined by the Bradford dye binding assay. Absorbance was
 EDTA. The protein content of the eluted fractions was
 20 tris/acetate buffer (pH 5.8) containing 50 mM NaCl and 1 mM
 (1.5 x 38 cm) (Pharmacia) pre-equilibrated with 5 mM bis-
 gel filtration chromatography using a G-25 Sephadex column
 stream of nitrogen gas. Excess unreacted 2-IT was removed by
 3 mM. The sample was incubated for 90 minutes at 4°C under a
 TEA/HCl (pH 8.0) was then added for a final concentration of
 15 volume was 2.8 ml. Seventeen uls of 2-imino-thiolane (2-IT)
 mM TEA/HCl and 28 ul 0.1 mM EDTA stock solutions. The final
 ethanolamine, pH 8.0) and 1 mM EDTA by addition of 300 ul 0.5
 (ddH₂O) was diluted to contain 60 mM TEA/HCl (tri-
 10 mg of avidin in 2.5 ml double distilled water
 Modification of Avidin - Egg White

EXAMPLE 2

conjugation.
 demonstrates that modified A108 was recovered after SPDP
 of SPDP-modified A108 is shown in Figure 1. Figure 1
 5 molecular weight A108 by gel filtration. The elution profile
 SPDP. The unreacted material was removed from the high
 through lysine and N-terminal amino acid modification using
 antibody (A108) through covalent coupling. A108 was modified
 FIGURE 1 shows the addition of SPDP to anti-EGFR

avidin were incubated together so that A108 would become covalently bound to the avidin through a sulphydryl linkage supplied by SPDP: 2-iminothiolane chemistry.

The immunocongugate composed of A108-avidin (labeled as conjugate in Figure 3) was separated from unreacted avidin by gel filtration. The peak of protein eluting into fractions 7-9 represents unmodified A108 and A108-avidin which was recovered and further purified.

EXAMPLE 4

Purification of Conjugate

Non-conjugated avidin was removed from the reaction mixture by gel filtration on a Pharmacia FPLC Superdex S-200 column (2.6 x 60 cm) pre-equilibrated with 20 mM Tris and 150 mM NaCl (pH 7.4). [FIGURE 3]

The antibody-avidin conjugate and free antibody fractions (6-10) were pooled and dialyzed (Spectra/Par molecular porous membrane tubing #2 MWCO 12,000-14,000) overnight against PBS at 4°C. The free antibody was removed from the mixture by use of a concavalin-A (Vector) agar se bound affinity. Column (1.5 cm x 7 cm) pre-equilibrated PBS (20 mM Na-K-phosphate, 150 mM NaCl, pH 7.0). After sample loading, the column was washed once with 40 ml of PBS containing 1 M NaCl (pH 7.0), and the conjugate was eluted with PBS containing 200 mM of methyl-D-mannose (pH 7.0) (fractions 34-38) (2 mlis each). The protein content of the eluted fractions was measured on a Varian Spectrophotometer at 280 nm.

Through binding of the carbohydrate moiety on avidin, A108-avidin was separated from A108 by its retention on an immobilized support of the plant lectin concavalin A (Con A) which binds alpha-methylmannoside (which is present on avidin). Free antibody has no alpha-methylmannoside and was washed through the Con A column (shown as free Ab on Figure 4). A108-avidin (labeled as conjugate) was displaced from the Con A-Sepharose column using elution with alpha-

35 chromatography on a 1.5 cm x 37 cm G-25 column equilibrated with PBS (pH 7.0). One ml fractions were collected in a

30 After 2 hours, the free biotin was separated by gel filtration. The sample was vortexed and incubated for 2 hours on ice. added to the gelonin in a clean, dry 13 x 100 mm glass tube. and immediately 20 μ l (0.2 mg) of this biotin solution was biotin was dissolved in 500 μ l dry dimethylformamide (DMF), gelonin in 2 ml of a 50 mM bicarbonate buffer (pH 8.5). 5 mg biotin to 1 mg gelonin) was used. The gelonin stock was 2 mg Co.). A five-fold molar excess of biotin to gelonin (= 0.1 mg succinimide ester long chain (NHS-LC) Biotin (Pierce Chemical Co.). The biotin used was in the form of N-hydroxy

25 Biotinylation of gelonin

EXAMPLE 5

gel filtration.

20 gelonin with NHS-biotin) and purified from unbound biotin by with biotin (by covalently bonding through lysine residues on gelonin). Purified gelonin protein was chemically modified when internalized into cells (e.g., the plant protein, utilizing a biotinylated protein having toxic activity only internalize into eukaryotic cells can be demonstrated

15 The ability of A108-avidin to bind biotin and was utilized in subsequent studies.

10 The sample applied to lane 7 (eluate from the Con A column) represents the purified immunconjugate (A108-avidin) which the polyacrylamide gel and destained to eliminate background. proteins were subjected to staining with Coomassie blue in conjugates of A108-avidin were resolved and visualized when shown in Figure 5, proteins representing A108, avidin or run to examine the purification steps of the conjugation. As their molecular size. A 7.5% acrylamide SDS-PAGE mini gel was electrophoresis (SDS-PAGE) which separates proteins based upon

5 The purity of the A108-avidin conjugate was monitored by sodium dodecyl sulfate polyacrylamide gel this procedure and was free of unmodified A108.

methylantranoside in solution. The conjugate was recovered by

35 A solution of gelonin in PBS was prepared in a concentration of 2 mg/ml. Next, a solution of biotinylated Tween-20.

30 The sample were then washed three times with PBS and 0.05% hours at room temperature with 5% bovine serum albumin in PBS. rinsed three times with PBS-0.05% Tween-20 and blocked for 1.5 overnight. Approximately 12 hours later, the samples were ng/well). The samples were covered and refrigerated of a Falcon 96-well microtiter plate was coated with 50 μ l (50 (1 μ g/ml solution). Using a multi-channel pipettor, each well coating buffer (50 mM NaHCO_3 (sodium bicarbonate, pH: 9.6) anti-gelonin antibody (10 μ l) (10 μ l) was diluted in 12 ml 25 A stock solution of 0.583 mg/ml murine monoclonal

Activity of biotinylated gelonin

EXAMPLE 6

20 proteins with an affinity for biotin.

15 incorporated into the gelonin molecule and recognized by assay. The results demonstrate that biotin can be increasing amounts of biotinylated gelonin placed into the retain biotin, based upon the increase in green color, by shown in Figure 6, the gelonin subjected to biotinylation did amount of biotin incorporated in the gelonin molecule. As amount of absorbance at 405 nm is directly proportional to the spectrophotometer at the wavelength of 405 nanometers. The substrate (ABTS) turns green and is measurable by peroxidase, which when incubated with a colorless horseradish 10 which was chemically conjugated to the enzyme horseradish was detected by rinsing the wells and adding streptavidin, anti-gelonin antibody. The retention of biotin with gelonin biotinylated gelonin were incubated in wells containing the gelonin protein. Indicated amounts of unmodified gelonin or polystyrene support with an antibody directed against the 5 gelonin, the biotinylated gelonin was immobilized on a To demonstrate that biotin was incorporated into the Bradford dye binding assay. Fractions 21-27 were pooled. Gilson fraction collector and assayed for protein content with

-11-

gelonin in PBS was prepared, also at concentration of 2 mg/ml.

100 μ l of 1 mg/ml solution of BSA in PBS was added to the plate, leaving the first row empty. To the first half of this

row, 200 μ l/well of the stock gelonin solution was added, and to the second half of the row 200 μ l/well of the biotinylated

stock solution was added. Using the multi-channel pipetor, 100 μ l was withdrawn from this first row and was mixed with

100 μ l BSA-PBS in the second row. This procedure was repeated from left to right across the plate resulting in the serial

dilution of the protein. The plate was covered and incubated for 1.5 hours at room temperature. The wells of the plate

were washed three times with PBS-0.05% Tween-20. 100 μ l of Avidin Peroxidase (Boehringer-Mannheim) diluted 1:6000 in

1 mg/ml BSA-PBS, was then added. Then the plate was incubated for 1.5 hours at room temperature followed by washing three

times with PBS-0.05% Tween-20. Finally, 100 μ l ABTS (2, 2'-amino-bis (3-ethyl benz Thiazoline - 6 - sulfonic acid))

hydrogen peroxide was added. The plate was read on a Biotek Laboratories Microplate Autoreader at 405 nm. [FIGURE 6B]

EXAMPLE 7

Conjugation of biotinylated gelonin with A108-avidin

A 5 molar excess of biotinylated gelonin to A108-

avidin was used. One ml (250 μ g) of A108-avidin was combined with 175 μ l (175 μ g) of biotinylated gelonin. The sample was

vortexed and incubated together for 1 hour at room temperature.

To remove unconjugated gelonin from the mixture, the

mixture was applied to a Pharmacia FPLC G-75 (1.6 x 60 cm) gel filtration column pre-equilibrated with 20 mM Tris containing

0.5 M NaCl (pH 7.4). One ml fractions were collected and read on a Varian Spectrophotometer at 280 nm. Peaks representing

A108-avidin biotinylated gelonin labeled conjugate is shown in Figure 7 and was free of unbound gelonin or free avidin

subunits.

The conjugation of A108-avidin with biotinylated

gelonin was examined using the same Elisa assay as described

Figure 9 shows the ability of the conjugate to transfer gelonin to the inside of the cell where it can induce cytotoxicity. The conjugate was incubated with cells that express EGF receptor at their cell surface (A431).

nm.

The plate was then read in the Microplate Autoreader at 540 nm.

Sorenson's buffer was added to extract dye from the cells. 20% methanol and rinsed in distilled water and 150 μ l of incubated 3 days and then stained with 0.5% crystal violet in (100 μ g/ml) antibody A108 added to each. The cells were same way but each with an addition of 100-fold molar excess conjugate. For a control, the conjugates were prepared in the plate, the final concentration was 1 μ g/ml for each to the plate. As there was already 100 μ l media present in prepared and 100 μ l of each dilution was added in triplicate (Corning). The A108-avidin:biotinylated gelonin was similarly filter, and serially diluted into ten 15 ml centrifuge tubes sterilized using a 0.22 micron Acrodisc (Gilmann) syringe prepare A108-avidin) in growing media was prepared, filter-linked to each other through the same chemistry used to using SPDP modified A108 and 2-IT modified gelonin covalently of A108-gelonin conjugate (Direct conjugate of A108-gelonin 37°C in a 5% CO₂ incubator. The next day, a 2 μ g/ml solution 96-well microtiter plate (Falcon) and incubated overnight at serum. 100 μ l of this solution was added to each well of a Laboratories) with 5% fetal bovine serum and 5% bovine calf acid, 100 mM glutamine and 50 μ l gentamicin (Tri-Bio media (MEM-minimum essential medium) with nonessential amino A431 cells were diluted 3 x 10⁴ cell/ml in growing media (MEM-minimum essential medium) with nonessential amino

EXAMPLE 8

5 biotinylated-gelonin. tested in the assay. Thus, the conjugate actually contains biotinylated gelonin or biotinylated gelonin:avidin-A108 was content could be detected by green color formation when either above for biotinylated gelonin. As shown in Figure 8, biotin

Internalization of the conjugate allows the intracellular effects of gelonin, i.e., cytotoxicity to occur. As shown in Figure 9, when conjugate was incubated with these cells, nanomolar (nM) concentrations (1×10^{-6} molar) killed A431 cells, demonstrating that A108-avidin can allow entry of biotinylated-gelonin into the A431 cell (open circles). When cells were co-incubated with free A108 antibody in a large excess (when compared to immunconjugate concentration) the ability of A108-avidin:biotinylated-gelonin to get into the A431 cell was impeded. Free A108 binds all the available EGF receptor on the cell surface, thus inhibiting immunconjugate from binding EGF receptor and introducing gelonin to the inside of the A431 cell. As shown in Figure 9, when A108 was present in 100-fold excess compared to the concentration of immunconjugate (closed circles), A431 cells were able to survive, demonstrating that the only way immunconjugate could enter and intoxicate cells was through its ability to bind and be internalized with EGF receptor. The direct covalent conjugate of A108-gelonin was also active in killing A431 by introduction of gelonin into the intracellular compartment of A431 cells (open triangles). Including 100-fold molar excess of free A108 with this direct conjugate also protected cells from intoxication by gelonin, demonstrating that the A108-gelonin was introduced to the intracellular compartment of the cell by its ability to interact and internalize with EGF receptor. Thus, utilization of the avidin:biotin interaction will allow introduction of a molecule into the intracellular compartment of the cell if directed and carried into that compartment with an antibody capable of recognizing an antigen on the cell which is internalized following engagement with the antigen (in this case the EGF receptor).

EXAMPLE 9

Effect of triple helix forming nucleic acid sequences on the expression of EGF receptor protein.

The ability of triple-helix forming oligonucleotide or nucleic acid sequences to suppress the expression of EGF

5 (which contains the same nucleic acids but in a random receptor gene promoter region or a non-sense control sequence nucleic acid sequence (#5 EGFR) capable of binding EGF were incubated for 72 hours with 40 μ M EGF receptor gene receptor protein in intact cell was demonstrated. A431 cells dishes containing 2×10^5 A431 cells. #5EGFR diluted in 2 ml media to a final concentration of 40 μ M was added to 1 dish. control nucleic acid sequences were diluted in 2 ml media to a final concentration of 40 μ M and this solution was added to the second dish of A431 cells. 2 ml of media only was added to the third dish. Incubation was for 72 hours. The cells were harvested by washing each dish three times with ice cold PBS. To solubilize the cell and extract protein, 1 ml RIPA buffer was added and the cells were released with a cell scraper and transferred to a centrifuge tube. Each tube was sonicated with a Kontes cell disrupter and centrifuged in a Sorvall ultra centrifuge for 1 hour at 4°C at 100,000 x g. After centrifugation, the supernatant was removed and protein content determined with the BCA protein assay (Pierce).

25 The EGF receptor in these extracts was immunoprecipitated with A108 antibody and the insoluble antibody-binding reagent pansorbin. Supernatant containing 200 μ g of protein was incubated for 2 hours at 4°C with 2.5 μ g of antibody A108. 50 μ l Pansorbin was added to each sample, vortexed and incubated for 30 minutes at 4°C. The EGF receptor bound to A108-pansorbin was washed to remove other proteins. After centrifugation for 1 minute (4°C) on a Sorvall microcentrifuge at 12,000 rpm, the supernatant was decanted. The pellet was washed by resuspending 3x with PBS containing 0.1% Triton followed by centrifugation and repelleting by centrifugation after each resuspension. After

final washing and centrifugation, the supernatant was poured off.

The EGF receptor present in each immunoprecipitate

was detected by incubating it with radioactive (^{32}P) labeled-ATP. The pellet was resuspended in 25 μl of 0.4 mM Na vanadate in 20 mM Hepes buffer (pH 7.4). Next, 25 μl of 20 mM Hepes buffer containing ^{32}P -labeled ATP (10 μCi) and 12 mM MnCl_2 was added. Samples were incubated for 5 minutes at room temperature. Since EGF receptor contains an enzyme activity capable of transferring ^{32}P from ATP to EGF receptor itself, the amount of ^{32}P transferred to EGF receptor becomes a measure of the quantity of EGF receptor in the immunoprecipitate from each of the lysates. The amount of radioactivity on EGF receptor can be compared by separating the EGF receptor from free ^{32}P ATP by SDS-PAGE and measuring radioactive EGF receptor by autoradiography of the polyacrylamide gel using commercial

x-ray film.

15 μl of 5X-Laemmli sample buffer was added to the sample, the sample was heated to 95°C for 5 minutes and then loaded onto a 7.5% polyacrylamide gel. The proteins were electrophoresed overnight at 14 mA. The gels were removed from the electrophoresis unit and fixed in 40% methanol, 10% acetic acid, 50% ddH_2O for 1 hour. The gel was dried on a Bio Rad 583 gel dryer for 2 hours at 80°C.

As can be seen in Figure 10, the incubation of cells with nucleic acid sequences directed against the EGF receptor gene promoter sequence (labeled anti-sense EGF) lowered the level of EGF receptor in A431 cells by five-fold when compared to random nucleic acid sequences (labeled non-sense EGF) or buffer alone (labeled control). Thus, incubation of cells with high concentrations of triple-helix forming nucleic acid sequences that interact with the promoter region of the EGF receptor gene suppress the expression of EGF receptor in intact A431 cells.

35 nucleic acid sequences are introduced into A431 cells through an avidin:biotin linkage. To demonstrate that nucleic acid sequences are directed against the EGF receptor gene promoter sequence is incorporated into A431 cells using A108 antibody, the A108-avidin chemically-linked conjugate is made and purified. Nucleic acid sequences are synthesized but with the substitution of a biotinylated-nucleotide for a normal nucleotide at one position in the sequence (preferably at the end or beginning of the sequence). These sequences are incubated with A108-avidin to allow complexes containing A108-avidin:biotin-nucleic acid sequences to form. These hybrid molecules are purified to remove free nucleic acid sequences and the amount of nucleic acid associated with A108-avidin is determined and is incubated with A431 cells. If gene expression-suppressing nucleic acid sequences are incorporated into the correct intracellular region of the A431 cells, it should be biochemically measurable. The specificity of the nucleic acid sequences to interrupt EGF receptor expression is determined by testing immunconjugates formed between anti-sense EGF, or non-sense EGF, with A108-avidin and measuring EGF receptor phosphorylation as described above in Example 9. In addition, the nucleic acid sequences' ability to get into the cell through formation of A108:EGF receptor complexes is tested by including a large molar excess of free A108 in the incubation mixture with A431 cells. The suppressive effects of hybrid molecules composed of antibody:nucleic acid sequences is inhibited in the presence of free A108 and confirms that nucleic acids are introduced into cells by their internalization with antibodies through engagement with an internalizable cell-surface antigen, e.g., the EGF receptor.

EXAMPLE 11

Larger biotinylated-nucleic acids are utilized to determine whether or not their entry into cells is mediated through the A108-avidin mechanism. The concentration of nucleic acid sequences presented to cells through the A108-

EXAMPLE 10

avidin mechanism which are necessary to alter EGF receptor levels will be compared to normal nucleic acid sequences free in solution. The ability of A108-avidin to direct active nucleic acid sequences to specific antigen expressing cells (e.g., the EGF receptor) is examined using cells which do not express this antigen. The delivery of these nucleic acids to specific cells within animals bearing cells which express this antigen is examined. Suppression of EGF receptor is measured. Both antisense or non-sense nucleic acid sequences complexed with A108-avidin are used (in addition to cells which do not express the specific antigen) as a measure of specificity of this delivery system and its intracellular biochemical specificity.

EXAMPLE 12

A monoclonal antibody against the breast carcinoma expressed antigen HER2/Neu (e.g., TAB 250) is chemically conjugated to avidin as described in Examples 1-4 for the A108 antibody. Antisense nucleic acid sequences against the antigen c-myc (5'-AACGTTGACGGCAT-3') are synthesized with a biotinylated adenine nucleotide replacing the adenosine at the terminal 5' position. The biotinylated adenine nucleic acid sequences are incubated with TAB-250-avidin and antibody-nucleic acid complexes composed of TAB-250:antisense c-myc are purified. These complexes are incubated with breast tumor cells expressing HER2/Neu, e.g., BT-474 cells or cells negative for this antigen, e.g., BT-20 cells. The suppression of c-myc by antisense nucleic acids is measured by western blotting for the c-myc protein from crude cell lysates. C-myc expression will be altered in HER2/Neu positive cells but not in antigen negative cells. Additionally, specificity is implied if a large excess (100-fold) of unmodified TAB-250 is able to inhibit the suppressive effects of the antibody-nucleic acid complexes on c-myc in antigen positive cell. The biotinylated nucleic acid sequence is altered by changing the position of the biotinylated-nucleotide within the sequence or by increasing the size of the sequence complementary to and

35 the c-Ha-ras oncogene. After incubation of immune:nucleic
carcinoma cells which express Lewis Y antigen and also contain
oligonucleotides are formed by incubation with T24 bladder
cytosine in the 5' position. BR96-antisense ras
with a biotinylated cytosine nucleotide in place of unmodified
30 mRNA sequence is chemically synthesized 5'-CAGCTGCAACCAAGC-3'
oligonucleotides complementary to the c-Ha-ras 5' flanking
conjugated to avidin as described above. Antisense
Lewis Y antigen on several human carcinomas is chemically
monoclonal antibody, BR96, which specifically binds

EXAMPLE 14

25 sequences through antigen internalization.
specificity and to confirm the mechanism of entry of antisense
coincubated with A108:nucleic acid complexes to assess
these target cells. As discussed above, excess A108
of growth suppression by preventing expression of bFGF in
20 conjugate, SNB cell growth is measured to determine the extent
promote their own growth. After incubation with this
critically dependent on the cells own synthesis of bFGF to
human glioma cells (SNB-19) which express EGF receptor and are
uncomplexed nucleic acid. These complexes are incubated with
15 antibody:nucleic acid complexes and are purified away from
sequences are incubated with A108-avidin to form
unmodified nucleotide at the 5'-terminal position. These
synthesized with a biotinylated guanosine in place of the
start site (5'- GGCTGCCATGCTCC- 3') are chemically
10 the basic fibroblast growth factor (bFGF) mRNA translation
oligonucleotides representing the complementary sequence to
chemically to avidin as described in Examples 1-4. Anti-sense
The monoclonal antibody A108 is conjugated

EXAMPLE 13

5 suppress c-myc expression.
breast carcinoma cells expressing HER2/Neu antigen which will
most specific and sensitive anti-sense sequence deliverable to
on the c-myc mRNA. The modifications are tested to obtain the
spanning the translation start site or first splice junction

acid complexes with T24 cells, the product of the ras oncogene, p21, is monitored by western blotting. Cell growth is also monitored. Neutralization of the effects of ras oncogene by intracellular delivery of antisense molecules through internalization of the Lewis Y antigen is demonstrated.

EXAMPLE 15

A108-avidin chemical conjugate is synthesized as described above and utilized to internalize synthetic double-stranded RNA molecules which are cytotoxic to specific carcinoma cells. A polymer of inosine (PI) is chemically synthesized (40mer) and hybridized to a polycytosine (PC)-39mer with a terminal cytosine derivatized with biotin. This double stranded RNA molecule is incubated with A108-avidin and the growth medium of ME-180 cervical carcinoma cells which express EGF receptor and are cytotoxicity sensitive to PI:PC (dsRNA). The intracellular delivery of dsRNA through internalization of the EGF receptor is measured by monitoring cell viability following incubation with this construct. Cytotoxicity induced by incubation with this conjugate will show that nucleic acids can be delivered to a specific set of cells with a non-viral vector.

All patents and publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference. One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The present examples along with the methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as

-20-

limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

5 WHAT IS CLAIMED IS:

Claims

1. A non-viral vector, comprising a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

2. The non-viral vector of claim 1, wherein said biotin-binding element is selected from the group consisting of avidin, streptavidin or analogues of avidin or streptavidin.

3. The non-viral vector of claim 1, wherein said cell binding element is a monoclonal antibody.

4. The non-viral vector of claim 3, wherein said monoclonal antibody specifically binds an antigen selected from the group consisting of epidermal growth factor receptor, c-erbB2 antigen, Lewis X antigen, transferrin receptor, MDR1, MDR3, insulin receptor, CD45, CD33, GP240, GD2, GD3, fibroblast growth factor receptor, platelet derived growth factor receptor.

5. The non-viral vector of claim 1, wherein said cell binding element is a ligand which specifically binds a cell surface receptor selected from the group consisting of transforming growth factor-alpha, heregulin, fibroblast growth factor, platelet-derived growth factor receptor.

6. The non-viral vector of claim 1, wherein said biotinylated moiety is selected from the group consisting of proteins and nucleic acids.

7. The non-viral vector of claim 6, wherein said protein is selected from the group consisting of gelonin, ricin, saporin, abrin, diphtheria toxin, pseudomonas exotoxin, ryalase, superoxide dismutase, protein tyrosine phosphatase,

protein phosphatase (PP-1 or PP-2), protein kinase A and protein kinase C.

8. The non-viral vector of claim 6, wherein said nucleic acid is selected from the group consisting of triplex forming oligonucleotides, anti-sense oligonucleotides, partial gene sequences and entire genes.

9. A method of introducing genetic material inside a specific cell comprising the administration of the non-viral vector to a human, wherein said non-viral vector comprises a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

10. The method of claim 9, wherein said biotin-binding element is selected from the group consisting of avidin, streptavidin or analogues of avidin or streptavidin.

11. The method of claim 10, wherein said cell binding element is a monoclonal antibody.

12. The method of claim 11, wherein said monoclonal antibody specifically binds an antigen selected from the group consisting of epidermal growth factor receptor, c-erbB2 antigen, Lewis X antigen, transferrin receptor, MDR1, MDR3, insulin receptor, CD45, CD33, BP240, GD2, GD3, fibroblast growth factor receptor, platelet derived growth factor receptor.

13. The method of claim 9, wherein said cell binding element is a ligand which specifically binds a cell surface receptor selected from the group consisting of transforming growth factor-alpha, heregulin, fibroblast growth factor, platelet-derived growth factor receptor.

14. The method of claim 9, wherein said biotinylated moiety is selected from the group consisting of proteins and nucleic acids.

15. The method of claim 14, wherein said protein is selected from the group consisting of gelonin, ricin, saporin, abrin, diphtheria toxin, pseudomonas exotoxin, ryalase, superoxide dismutase, protein tyrosine phosphatase, protein phosphatase (PP-1 or PP-2), protein kinase A and protein kinase C.

16. The method of claim 14, wherein said nucleic acid is selected from the group consisting of triplex forming oligonucleotides, anti-sense oligonucleotides, partial gene sequences and entire genes.

17. A method of delivering a cytotoxic moiety to a cell comprising the administration of a non-viral vector to a human, wherein said non-viral vector comprises a cell binding component having a biotin-binding element conjugated to a biotinylated moiety.

18. The method of claim 17, wherein said biotin-binding element is selected from the group consisting of avidin, streptavidin or analogues of avidin or streptavidin.

19. The method of claim 18, wherein said cell binding element is a monoclonal antibody.

20. The method of claim 17, wherein said monoclonal antibody specifically binds an antigen selected from the group consisting of epidermal growth factor receptor, c-erbB2 antigen, Lewis X antigen, transferrin receptor, MDR1, MDR3, insulin receptor, CD45, CD33, GP240, GD2, GD3, fibroblast growth factor receptor, platelet derived growth factor receptor.

-24-

21. The method of claim 17, wherein said cell binding element is a ligand which specifically binds a cell surface receptor selected from the group consisting of transforming growth factor- α , heregulin, fibroblast growth factor, platelet-derived growth factor receptor.

22. The method of claim 17, wherein said biotinylated moiety is selected from the group consisting of proteins and nucleic acids.

23. The method of claim 22, wherein said protein is selected from the group consisting of gelonin, ricin, saporin, abrin, diphtheria toxin, pseudomonas exotoxin, ryalase, superoxide dismutase, protein tyrosine phosphatase, protein phosphatase (PP-1 or PP-2), protein kinase A and protein kinase C.

24. The method of claim 22, wherein said nucleic acid is selected from the group consisting of triplex forming oligonucleotides, anti-sense oligonucleotides, partial gene sequences and entire genes.




FIG. 1

1 / 11




FIG. 2




FIG. 3





FIG. 4

FIG. 5






FIG. 6A




FIG. 6B

7/11




FIG. 7

9/11




FIG. 8





FIG.9

SUBSTITUTE SHEET (RULE 26)

FIG. 10

Control	AS	Sense	(Anti-sense)
---------	----	-------	--------------



11/11

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/01161

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07K 16/00, 16/46, 19/00; A61K 48/00, 51/00, 121/00
US CL : 530/391.1, 350; 424/178.1, 179.1, 181.1, 182.1, 183.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/391.1, 350; 424/178.1, 179.1, 181.1, 182.1, 183.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS/DIALOG, EMBASE, MEDLINE, BIOSYS, LIFESCI, WPI

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
-----------	--	-----------------------

Y	US, A, 5,026,785 (MAGE ET AL.) 25 JUNE 1991, ENTIRE DOCUMENT.	1-24
Y	TRENDS IN BIOTECHNOLOGY, VOL. 10, ISSUED AUGUST 1992, E. WICKSTROM, "STRATEGIES FOR ADMINISTERING TARGETED THERAPEUTIC OLIGODEOXYNUCLEOTIDES", PAGES 281-287, SEE ENTIRE DOCUMENT.	1-24
Y	BIOTECHNOLOGY, VOL. 9, ISSUED DECEMBER 1991, T. SANO ET AL., "A STREPTAVIDIN-PROTEIN A CHIMERA THAT ALLOWS ONE-STEP PRODUCTION OF A VARIETY OF SPECIFIC ANTIBODY CONJUGATES", PAGES 1378-1381, SEE ENTIRE DOCUMENT.	1-24
Y	EP, A, 0,496,074 (PAGANELLI ET AL.) 29 JULY 1992, SEE ENTIRE DOCUMENT.	1-24

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:	T	X	Y	G
A* document defining the general state of the art which is not considered to be of particular relevance				
E* earlier document published on or after the international filing date				
L* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)				
O* document referring to an oral disclosure, use, exhibition or other means				
P* document published prior to the international filing date but later than the priority date claimed				

Date of the actual completion of the international search	09 MAY 1995
Date of mailing of the international search report	22 MAY 1995

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <i>Christopher E. Eischenschek</i> CHRISTOPHER EISENSCHENK Telephone No. (703) 308-0196
--	---